# TOBACCO FACT SHEET. Youth Smoking

Cigarette smoking is the chief preventable cause of death and disease in the United States, accounting for more than one in every five deaths. Even in light of the evidence, more than one million U.S. teenagers begin smoking each year. Although high school students experiment with cigarette smoking at similar rates, those living in larger cities are less likely to continue use than are their peers who live in smaller cities and towns.



## Mississippi Facts

<b>_</b>	Approximately 60 Mississippi youth begin smoking every day.
	28 percent of high school students are regular cigarette smokers.

- ☐ 76 percent of high school students have tried smoking cigarettes, meaning one or two puffs.
- □ 58 percent of Mississippi youth have smoked an entire cigarette at least once. Among these, 48 percent were age 13 or younger when this event first occurred.

#### A Healthy People 2000 National Objective:

■ Smoking cigarettes can cause impotence.

To reduce the initiation of cigarette smoking by children and youth so that no more than 15 percent have become regular cigarette smokers by age 20.

### The Facts

THE Lacts					
	89 percent of adult daily smokers begin before the age of 18.				
	Every day, 3,000 young people become regular smokers in the United States.				
	Tobacco is often the first drug - gateway drug - used by youth who later use other drugs like alcohol and marijuana.				
	Youth who start smoking are more likely to get lower grades in school, to hang out with other kids who smoke, to have a low self-image, to become involved in acts of violence, and don't know how to say no to tobacco.				
	Cigarette advertisements are designed to make youth think that smoking is cool and that everyone does it. These misleading ads appear to increase youth desire to smoke.				
	85 percent of youth who smoke, use the 3 brands most publicized: Marlboro, Camel and Newport. Only 35 percent of adults smoke these brands.				

# The Risks

	The health effects of smoking build up over time. Smokers are at a higher risk than non-smokers for					
	Addiction to nicotine		Diabetes			
	Heart attack		Emphysema			
	Stroke		Miscarrage			
	High blood pressure		Stillborn			
	Lung cancer		or low birthweight babies			
Immediate consequences for cigarette smokers are						
	Bad breath		Smellyclothes			
	Yellow teeth		Premature wrinkles and aging			
	Coughing		Phlegm production			
	Dizziness		Nausea			
	Wheezing		Shortness of breath			
What Can Be Done  By Adults:  Set a good example. If you have children, they are three times as likely to become smokers						
	if you smoke.  Work to ban cigarette smoking in schools, restaurants, and other public places					
	where children frequent.  If you smoke, stop now. It not only helps you, but also your children.					
By Youth:  ☐ Don't start. ☐ Encourage peers not to start. ☐ Educate yourself about the harmful effects of tobacco use. The typical smoker spends about \$700 a year on cigarettes.						
			Produced by — Don't Start			



Mississippi State Department of Health Post Office Box 1700 2423 North State Street

Jackson, Mississippi 39215-1700

**Equal Opportunity in Employment Services** 

This program is funded in whole or in part with federal money; the FY 1995 appropriation from tHe Preventive Health and Health Services Block Grant is \$2,363,845.